



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/653 // (A01N 43/653, 43:36, 43:30, 43:54)	A1	(11) International Publication Number: WO 98/57543 (43) International Publication Date: 23 December 1998 (23.12.98)						
<p>(21) International Application Number: PCT/EP98/03571</p> <p>(22) International Filing Date: 12 June 1998 (12.06.98)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>9712466.3</td> <td>16 June 1997 (16.06.97)</td> <td>GB</td> </tr> <tr> <td>9712463.0</td> <td>16 June 1997 (16.06.97)</td> <td>GB</td> </tr> </table> <p>(71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).</p> <p>(71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1235 Wien (AT).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): MÜLLER, Kaspar [CH/CH]; Niederfeldweg 2, CH-4124 Schönenbuch (CH). KNAUF-BEITER, Gertrude [DE/DE]; Fritz-Fischerstrasse 12, D-79379 Müllheim (DE). STECK, Bernhard [DE/CH]; Im Marcoup 9, CH-3286 Muntelier (CH).</p> <p>(74) Agent: BECKER, Konrad; Novartis AG, Patent- und Markenabteilung, Lichtstrasse 35, CH-4002 Basel (CH).</p>		9712466.3	16 June 1997 (16.06.97)	GB	9712463.0	16 June 1997 (16.06.97)	GB	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published</p> <p><i>With international search report.</i></p> <p><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
9712466.3	16 June 1997 (16.06.97)	GB						
9712463.0	16 June 1997 (16.06.97)	GB						
<p>(54) Title: PESTICIDAL COMPOSITIONS</p>								
<p>(57) Abstract</p> <p>Fungicidal mixtures having synergistically enhanced action against disease infestation in plants, comprising at least two active ingredient components together with a suitable carrier, wherein component I is: I) fludioxonil (=4-(2,2-difluoro-1,3-benzodioxol-4-yl)pyrrole-3-carbonitrile) and wherein component (II) is: IIA) triticonazole (=5-[(4-chlorophenyl)methylene]-2,2-dimethyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol), or IIB) cyproconazole (=α-(4-chlorophenyl)-α-(1-cyclopropylethyl)-1-H-1,2,4-triazol-1-ethanol). These mixtures are especially useful for seed treatment.</p>								

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Pesticidal compositions

The present invention relates to fungicidal mixtures having synergistically enhanced action and to methods of using such mixtures in crop protection, especially in the control and prevention of disease infestation on seed.

The mixtures according to the invention comprise at least two fungicidally active components together with a suitable carrier material, wherein component I is
I) fludioxonil (= 4-(2,2-difluoro-1,3-benzodioxol-4-yl)pyrrole-3-carbonitrile) (The Pesticide Manual, 10th. edition, 1994, 326);
and wherein component II is
IIA) triticonazole (= 5-[(4-chlorophenyl)methylene]-2,2-dimethyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol) (The Pesticide Manual, 10th. edition, 1994, 712) or
IIB) cyproconazole (= α -(4-chlorophenyl)- α -(1-cyclopropylethyl)-1-H-1,2,4-triazol-1-ethanol) (The Pesticide Manual, 10th. edition, 1994, 183).

The active ingredient mixtures I+II according to the invention have very advantageous properties for protecting plants against disease infestation, especially in the control and prevention of disease infestation on seed.

The synergistically enhanced action of mixtures of components I and II manifests itself, for example, in lower disease infestation, lower rates of application, a longer duration of action and altogether higher crop yields. Such enhancements were not to be expected from the sum of the actions of the individual components.

The weight ratio is selected as to give a synergistic action. In general, the weight ratio I:II is from 100:1 to 1:100.

Advantageous mixing ratios of the two active ingredients are

I:IIA = 1:20 to 10:1, preferably 1:15 to 1:1, most preferably 1:10 to 1:2.

I:IIB = 1:10 to 20:1, preferably 1:5 to 10:1, most preferably 1:1 to 5:1.

The above mentioned mixtures may be mixed with other pesticides, preferably fungicides as for example cyprodinil. Particularly preferred is the 3-component mixture comprising

I) fludioxonil,

IIb) cyproconazole and

III) cyprodinil (= 4-cyclopropyl-6-methyl-N-phenyl-2-pyrimidinamin (The Pesticide Manual, 10th. edition, 1994, 109). In this case advantageous mixing ratios are

I:IIb:III= 1:10:40 to 20:1:10, preferably 1:5:20 to 10:1:10, most preferably 1:1:4 to 5:1:10.

Further advantageous synergistic mixtures are

triticonazole + cyprodinil, cyproconazole + R-metalaxyl, and fludioxonil + R-metalaxyl + flutriafol.

The active ingredient mixtures in question can be used to inhibit or destroy the microorganisms which occur on plants or on parts of plants (the fruit, blossom, leaves, stems, tubers or roots) of different crops of useful plants, while at the same time parts of plants that grow later are also protected against such microorganisms. They can also be used as dressings in the treatment of plant propagation material, especially seed (fruit, tubers, grains) and plant cuttings (e.g. rice), to provide protection against fungus infections as well as against phytopathogenic fungi which occur in the soil. The active ingredient mixtures according to the invention are distinguished by the fact that they are especially well tolerated by plants and are environmentally friendly.

The active ingredient mixtures are effective against phytopathogenic fungi belonging to the following classes: Ascomycetes (e.g. Venturia, Podosphaera, Erysiphe, Monilinia, Mycosphaerella, Uncinula); Basidiomycetes (e.g. the genus Hemileia, Rhizoctonia, Puccinia); Fungi imperfecti (e.g. Botrytis, Helminthosporium, Rhynchosporium, Fusarium, Septoria, Cercospora, Alternaria, Pyricularia and Pseudocercospora herpotrichoides); Oomycetes (e.g. Phytophthora, Peronospora, Bremia, Pythium, Plasmopara).

Target crops for the areas of indication disclosed herein comprise within the scope of this invention e.g. the following species of plants: cereals (wheat, barley, rye, oats, rice, sorghum and related crops); beet (sugar beet and fodder beet); pomes, stone fruit and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries); leguminous plants (beans, lentils, peas, soybeans); oil plants (rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans, groundnuts); cucumber plants (marrows, cucumbers, melons); fibre plants (cotton, flax, hemp, jute); citrus fruit

(oranges, lemons, grapefruit, mandarins); vegetables (spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, paprika); lauraceae (avocados, cinnamon, camphor); or plants such as maize, tobacco, nuts, coffee, sugar cane, tea, vines, hops, bananas and natural rubber plants, as well as ornamentals (flowers, shrubs, broad-leaved trees and evergreens, such as conifers). This list does not represent any limitation.

The active ingredient mixtures according to the invention are especially advantageous for seed treatment of cereals (wheat, barley, rye, oats, rice, sorghum and related crops), in particular seed of wheat and barley.

The mixtures of compounds of formulae I and II are normally used in the form of compositions. The compounds of formulae I and II can be applied to the area or plant to be treated either simultaneously or in succession on the same day, if desired together with further carriers, surfactants or other application-promoting adjuvants customarily employed in formulation technology.

Suitable carriers and adjuvants can be solid or liquid and are the substances ordinarily employed in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, thickeners, binders or fertilisers.

A method of applying an active ingredient mixture comprising at least one of each of the active ingredients I and II is application to the parts of the plants that are above the soil, especially to the leaves (foliar application). The frequency and rate of application depend upon the biological and climatic living conditions of the pathogen. The active ingredients can, however, also penetrate the plant through the roots *via* the soil or *via* the water (systemic action) if the locus of the plant is impregnated with a liquid formulation (e.g. in rice culture) or if the substances are introduced in solid form into the soil, e.g. in the form of granules (soil application). In order to treat seed, the compounds of formulae I and II can also be applied to the seeds (coating), either by impregnating the tubers or grains with a liquid formulation of each of the active ingredients in succession, or by coating them with an already combined wet or dry formulation. In addition, in special cases, other methods of application to plants are possible, for example treatment directed at the buds or the fruit trusses.

The compounds of the combination are used in unmodified form or, preferably, together with the adjuvants conventionally employed in formulation technology, and are therefore formulated in known manner e.g. into emulsifiable concentrates, coatable pastes, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granules, or by encapsulation in e.g. polymer substances. As with the nature of the compositions, the methods of application, such as spraying, atomising, dusting, scattering, coating or pouring, are chosen in accordance with the intended objectives and the prevailing circumstances. Advantageous rates of application of the active ingredient mixture are generally from 50 g to 2 kg a.i./ha, especially from 100 g to 1000 g a.i./ha, more especially from 250 g to 700 g a.i./ha. In the case of the treatment of seed, the rates of application are from 0.5 g to 500 g, preferably from 1 g to 100 g, most preferably from 5 g to 50 g a.i. per 100 kg of seed.

The formulations are prepared in known manner, e.g. by homogeneously mixing and/or grinding the active ingredients with extenders, e.g. solvents, solid carriers and, where appropriate, surface-active compounds (surfactants).

Suitable solvents are: aromatic hydrocarbons, preferably the fractions containing 8 to 12 carbon atoms, e.g. xylene mixtures or substituted naphthalenes, phthalates, such as dibutyl phthalate or dioctyl phthalate, aliphatic hydrocarbons, such as cyclohexane or paraffins, alcohols and glycols and their ethers and esters, such as ethanol, ethylene glycol, ethylene glycol monomethyl or monoethyl ether, ketones, such as cyclohexanone, strongly polar solvents, such as N-methyl-2-pyrrolidone, dimethyl sulfoxide or dimethylformamide, as well as vegetable oils or epoxidised vegetable oils, such as epoxidised coconut oil or soybean oil; or water.

The solid carriers used, e.g. for dusts and dispersible powders, are normally natural mineral fillers, such as calcite, talcum, kaolin, montmorillonite or attapulgite. In order to improve the physical properties it is also possible to add highly dispersed silicic acid or highly dispersed absorbent polymers. Suitable granulated adsorptive carriers are porous types, for example pumice, broken brick, sepiolite or bentonite, and suitable nonsorbent carriers are, for example, calcite or sand. In addition, a great number of pregranulated materials of inorganic or organic nature can be used, e.g. especially dolomite or pulverised plant residues.

Depending upon the nature of the compounds of formulae I and II to be formulated, suitable surface-active compounds are non-ionic, cationic and/or anionic surfactants having good emulsifying, dispersing and wetting properties. The term "surfactants" will also be understood as comprising mixtures of surfactants.

Particularly advantageous application-promoting adjuvants are also natural or synthetic phospholipids of the cephalin and lecithin series, e.g. phosphatidylethanolamine, phosphatidylserine, phosphatidylglycerol and lysolecithin.

The agrochemical compositions normally comprise 0.1 to 99 %, especially 0.1 to 95 %, compounds of formulae I and II, 99.9 to 1 %, especially 99.9 to 5 %, of a solid or liquid adjuvant and 0 to 25 %, especially 0.1 to 25 %, of a surfactant.

Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations.

The Examples which follow serve to illustrate the invention, "active ingredient" denoting a mixture of compound I and compound II in a specific mixing ratio.

Formulation Examples

<u>Wettable powders</u>	a)	b)	c)
active ingredient [I:II = 1:3(a), 1:2(b), 1:1(c)]	25 %	50 %	75 %
sodium lignosulfonate	5 %	5 %	-
sodium lauryl sulfate	3 %	-	5 %
sodium diisobutyl-naphthalenesulfonate	-	6 %	10 %
phenol polyethylene glycol ether (7-8 mol of ethylene oxide)	-	2 %	-
highly dispersed silicic acid	5 %	10 %	10 %
kaolin	62 %	27 %	-

The active ingredient is thoroughly mixed with the adjuvants and the mixture is thoroughly ground in a suitable mill, affording wettable powders which can be diluted with water to give suspensions of the desired concentration.

- 6 -

<u>Dusts</u>	a)	b)	c)
active ingredient [I:II = 1:6(a), 1:2(b), 1:10(c)]	5 %	6 %	4 %
talcum	95 %	-	-
kaolin	-	94 %	-
mineral filler	-	-	96 %

Ready-for-use dusts are obtained by mixing the active ingredient with the carrier and grinding the mixture in a suitable mill. Such powders can also be used for dry dressings for seed.

Coated granules

active ingredient (I:II = 1:10)	8 %
polyethylene glycol (mol. wt. 200)	3 %
kaolin	89 %

The finely ground active ingredient is uniformly applied, in a mixer, to the kaolin moistened with polyethylene glycol. Non-dusty coated granules are obtained in this manner.

Suspension concentrate

active ingredient (I:II = 1:8)	40 %
propylene glycol	10 %
nonylphenol polyethylene glycol ether (15 mol of ethylene oxide)	6 %
sodium lignosulfonate	10 %
carboxymethylcellulose	1 %
silicone oil (in the form of a 75 % emulsion in water)	1 %
water	32 %

The finely ground active ingredient is intimately mixed with the adjuvants, giving a suspension concentrate from which suspensions of any desired dilution can be obtained by dilution with water. Using such dilutions, living plants as well as plant propagation material can be treated and protected against infestation by microorganisms, by spraying, pouring or immersion.

Biological Examples

A synergistic effect exists whenever the action of an active ingredient combination is greater than the sum of the actions of the individual components.

The action to be expected E for a given active ingredient combination obeys the so-called COLBY formula and can be calculated as follows (COLBY, S.R. "Calculating synergistic and antagonistic responses of herbicide combination". Weeds, Vol. 15, pages 20-22; 1967):

ppm = milligrams of active ingredient (= a.i.) per litre of spray mixture

X = % action by active ingredient I using p ppm of active ingredient

Y = % action by active ingredient II using q ppm of active ingredient.

According to Colby, the expected (additive) action of active ingredients I+II using p+q ppm

of active ingredient is $E = X + Y - \frac{X \cdot Y}{100}$

If the action actually observed (O) is greater than the expected action (E), then the action of the combination is superadditive, i.e. there is a synergistic effect.

Example B-1: Action against Drechslera teres on barley

10-day-old barley plants of the "Golden Promise" variety are sprayed with a spray mixture of the active ingredient mixture. The treated plants are infected 24 hours later with a conidia suspension of the fungus and incubated in a climatic chamber at 70 % relative humidity and 20-22°C. Fungus infestation is evaluated 5 days after infection. The mixtures according to the invention exhibit good synergistic actions.

Example B-2: Action against Helminthosporium gramineum on barley; seed treatment

Barley seeds are contaminated with a spore suspension of the fungus and left to dry. The contaminated seeds are dressed with a suspension of the test substance. After two days, the seeds are placed on suitable agar dishes and, after a further four days, the development of the fungal colonies around the seeds is evaluated. The number and size of the fungal colonies are used to evaluate the test substance.

Results

component I: Fludioxonil

component IIA: Triticonazole

Test no.	g a.i. per 100 kg seed		I:II	% action		SF O/E
	a.i. I	a.i. IIA		O (observed)	E (expected)	
1	0.5			23		
2		10		5		
3	0.5	10	1:20	41	27	1.5

Example B-3: Action against Fusarium nivale in rye; seed treatment

Rye seeds of the Tetrahell variety naturally infected with *Fusarium nivale* are dressed in a roller mixer with the test fungicides. The infected and treated rye is sown in October in the open with a seeder in plots 3 metres long and in 6 rows. Three replicates are carried out with each concentration. The test plants are cultivated under normal field conditions, harvested and dried.

Results

component I: Fludioxonil

component IIB: Cyproconazole

Test no.	g a.i. per 100 kg seed		I:II	dry weight of crop (O: observed; E: expected)			SF
	a.i. I	a.i. IIB		O [kg]	% increase over check		O/E
					O	E	
0 (check)	-	-		767			
1	0.5			770	0		
2		1		880	15		
3	0.5	1	1:2	973	27	15	1.8


Claims

1. A composition having synergistic action against disease infestation in plants, comprising at least two active ingredient components together with a suitable carrier, wherein component I is
I) fludioxonil (= 4-(2,2-difluoro-1,3-benzodioxol-4-yl)pyrrole-3-carbonitrile)
and wherein component II is
IIA) triticonazole (= 5-[(4-chlorophenyl)methylene]-2,2-dimethyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol) or
IIB) cyproconazole (= α -(4-chlorophenyl)- α -(1-cyclopropylethyl)-1-H-1,2,4-triazol-1-ethanol).
 2. A composition according to claim 1, wherein component II is IIA) triticonazole.
 3. A composition according to claim 1, wherein component II is IIB) cyproconazole.
 4. A composition according to claim 1, wherein the ratio by weight of I:IIA is 1:20 to 10:1.
 5. A composition according to claim 1, wherein the ratio by weight of I:IIB is 1:10 to 20:1.
 6. A composition according to claim 3, which additionally comprises III) cyprodinil.
 7. A method of protecting plants against plant diseases by treating the plants, parts of plants or their surroundings with a component I and a component II according to claim 1, in any desired sequence or simultaneously.
 8. A method according to claim 7, wherein seed is treated.
 9. A method according to claim 8, wherein seed of cereals is treated.
 10. Seed that has been treated in accordance with claim 8.
-

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 98/03571

A. CLASSIFICATION OF SUBJECT MATTER		
IPC6: A01N 43/653 // (A01N 43/653, 43:36, 43:30, 43:54) According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
IPC6: A01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
CA, WPI		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	GB 2308301 A (RHONE-POULENC AGROCHIMIE), 25 June 1997 (25.06.97) --	1-10
A	EP 0487451 A1 (CIBA-GEIGY AG), 27 May 1992 (27.05.92) --	1-10
A	EP 0513567 A1 (BAYER AG), 19 November 1992 (19.11.92) --	1-10
A	EP 0575838 A1 (RHONE-POULENC AGROCHIMIE), 29 December 1993 (29.12.93), claims 1,2,9 --	1-10
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
20 October 1998		25. 11. 98
Name and mailing address of the ISA:  European Patent Office, P.B. 5818 Patentlaan 2 NL-2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer GERD STRANDELL

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 98/03571

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0467792 A1 (RHONE-POULENC AGROCHIMIE), 22 January 1992 (22.01.92), page 4, line 45 - line 46, the claims --	1-10
A	EP 0630569 A2 (CIBA-GEIGY AG), 28 December 1994 (28.12.94), page 7, line 4 - line 44, the claims --	1-10
A	EP 0548025 A1 (CIBA-GEIGY AG), 23 June 1993 (23.06.93) -- -----	1-10

INTERNATIONAL SEARCH REPORT
Information on patent family members

27/07/98

International application No.
PCT/EP 98/03571

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
GB	2308301	A	25/06/97	FR	2742311 A,B	20/06/97
				GB	9626306 D	00/00/00

EP	0487451	A1	27/05/92	AT	136424 T	15/04/96
				AU	639841 B	05/08/93
				AU	8790691 A	21/05/92
				BG	60133 A	15/11/93
				CA	2055529 A	17/05/92
				CS	9103468 A	17/06/92
				DE	59107663 D	00/00/00
				DK	487451 T	06/05/96
				ES	2085455 T	01/06/96
				FI	915362 A	17/05/92
				GR	3019631 T	31/07/96
				HR	950140 A	31/12/96
				HU	209690 B	28/10/94
				IE	72098 B	12/03/97
				IL	100051 A	18/06/96
				JP	2646164 B	25/08/97
				JP	4266804 A	22/09/92
				LT	1504 A	26/06/95
				LT	3727 B	26/02/96
				LV	10993 A,B	20/02/96
				MD	277 B	30/09/95
				NO	180405 B,C	06/01/97
				PL	167802 B	30/11/95
				PT	99517 A	30/09/92
				RU	2040899 C	09/08/95
				SG	46991 A	20/03/98
				US	5250556 A	05/10/93

EP	0513567	A1	19/11/92	AT	130162 T	15/12/95
				AU	648404 B	21/04/94
				AU	1515692 A	05/11/92
				CA	2067644 A	04/11/92
				DE	4114447 A	05/11/92
				DE	59204301 D	00/00/00
				DK	513567 T	25/03/96
				ES	2079093 T	01/01/96
				GR	3018099 T	29/02/96
				HU	212606 B	30/09/96
				JP	5148110 A	15/06/93
				MX	9201842 A	01/11/92
				PL	168731 B	29/03/96
				RU	2042327 C	27/08/95
				US	RE34983 E	27/06/95
				US	5198456 A	30/03/93

INTERNATIONAL SEARCH REPORT

Information on patent family members

27/07/98

International application No.

PCT/EP 98/03571

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0575838 A1	29/12/93	AT 138243 T DE 69302756 D,T DK 575838 T ES 2087612 T GR 3020006 T JP 6056605 A	15/06/96 02/10/96 05/08/96 16/07/96 31/08/96 01/03/94
EP 0467792 A1	22/01/92	AT 122205 T AU 656320 B AU 7834091 A CA 2043812 A CN 1063396 A CS 9101784 A DE 69109560 D,T DK 467792 T EG 19259 A ES 2071958 T FI 97195 B,C FI 912822 A FR 2663196 A HU 209725 B IE 67894 B IL 98400 A JP 4230204 A OA 9364 A PT 97934 A RU 2035140 C TR 25435 A US 5290791 A US 5629330 A	15/05/95 02/02/95 19/12/91 14/12/91 12/08/92 15/01/92 28/09/95 10/07/95 30/11/94 01/07/95 31/07/96 14/12/91 20/12/91 28/10/94 01/05/96 10/03/98 19/08/92 15/09/92 30/04/92 20/05/95 01/03/93 01/03/94 13/05/97
EP 0630569 A2	28/12/94	AT 168856 T AU 677392 B AU 6599094 A BG 98869 A BR 9402556 A CA 2126714 A CN 1099220 A CZ 9401564 A EP 0818144 A EP 0818145 A HU 67004 A HU 9401932 D IL 110132 A JP 7017811 A NZ 260840 A PL 303960 A SI 9400265 A SK 76794 A US 5430035 A US 5536726 A	15/08/98 24/04/97 05/01/95 31/03/95 28/03/95 29/12/94 01/03/95 18/01/95 14/01/98 14/01/98 30/01/95 00/00/00 05/04/98 20/01/95 27/11/95 09/01/95 31/12/94 05/01/95 04/07/95 16/07/96

INTERNATIONAL SEARCH REPORT

Information on patent family members

27/07/98

International application No.

PCT/EP 98/03571

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0548025 A1	23/06/93	AT 136728 T	15/05/96
		AU 652302 B	18/08/94
		AU 3025892 A	24/06/93
		BR 9205070 A	22/06/93
		CA 2085714 A	20/06/93
		DE 59206041 D	00/00/00
		DK 548025 T	13/05/96
		ES 2086716 T	01/07/96
		GR 3019667 T	31/07/96
		HU 212479 B	29/07/96
		IL 104116 A	18/02/97
		JP 2753932 B	20/05/98
		JP 5271012 A	19/10/93
		MX 9207454 A	01/06/93
		NZ 245508 A	26/09/95
		PL 173377 B	27/02/98
		RU 2098962 C	20/12/97
		SG 47444 A	17/04/98
		SK 374792 A	10/08/94
		TR 26346 A	15/03/95
		US 5403844 A	04/04/95
		US 5567705 A	22/10/96
		US 5627188 A	06/05/97
		ZA 9209824 A	21/06/93
